

*Evaluation of Interleukin-2 to Detect
Active and Latent Tuberculosis among
Household Contacts of Pulmonary
Tuberculosis Cases*

By : Dr. Azadeh Jafarsteh



BACKGROUND



- ✘ The interferon-gamma release assays (IGRAs) are the most important diagnostic approach to *Mycobacterium tuberculosis* infection diagnosis. However, they cannot discriminate between latent tuberculosis infection (LTBI) and active tuberculosis (TB). Some recent studies suggested that interleukin-2 (IL-2) response to *M. tuberculosis* could be utilized as a potential biomarker to discriminate active disease from LTBI.

OBJECTIVES



- ✘ The present study aimed at evaluating the diagnostic potential of IL-2 to identify active TB and LTBI in household contacts of patients with pulmonary TB in two TBpandemic provinces of Iran. Additionally, the IL-2 assay findings were compared with those obtained from QFTPlus to assume whether the diagnostic performance of IGRA could be improved by combining it with the IL-2 assay

× METHODS



STUDY DESIGN AND POPULATION



- ✘ The current study was conducted in various districts of two high TB burden provinces in Iran, including Golestan and Sistan-Baluchestan. The current study protocol was approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences (ethical code: IR.NIMAD.REC.1395.039). Informed written consent was taken from all the patients prior to enrollment in the study. From August 2017 to August 2019, a total of 650 household contacts of patients with pulmonary TB were invited to participate in the study if they met the definition of a household contact..

STUDY DESIGN AND POPULATION

- ✘ The inclusion criteria were: children aged < 18 years, with known contact with an index TB case within the previous six months, and households with close contacts with an index TB case. HIV-positive patients were excluded from the study. Before blood sampling, all participants signed their informed consent form.

. All subjects were diagnosed on clinical, radiological, and laboratory findings, including TB contact history, chest X-ray radiography, clinical sign\symptoms, microscopic examination of sputum smear, and bacterial culture test for *M. tuberculosis*. LTBI status was defined as a positive IGRA reaction without clinical/radiological evidence of active TB disease

QFT-PLUS ASSAY



- ✘ The QFT-Plus test was performed according to the manufacturer's instructions (QIAGEN, Germantown, MD, USA). Briefly, 1 mL of whole blood was added to QFT-Plus tubes, including nil control tube, TB1 tube (comprising mycobacterial antigens derived from ESAT-6 and CFP-10 proteins of *M. tuberculosis* stimulating CD4 T-cells), TB2 tube (containing *M. tuberculosis* antigens that stimulate CD8 T-cells in addition to CD4 T-cells), and mitogen control tube.

QFT-PLUS ASSAY

- × . All tubes were incubated at 37 °C for 20 hours; then, the samples were centrifuged, and the plasma supernatants were analyzed to measure the concentration of IFN- γ by ELISA technique. QFT-Plus tests were considered positive if the antigen-stimulated response of IFN- γ (TBAg-*Nil*) was 0.35 IU/mL, negative if the mitogen-stimulated response (Mitogen-*Nil*) was 0.5 IU/ml and the antigen-stimulated response was 0.35 IU/mL, or indeterminate if both mitogenstimulated and antigen-stimulated responses were 0.35 IU/mL and unstimulated response (*Nil*) was 0.8 IU/mL

IL-2 MEASUREMENT



- ✘ plasma levels of IL-2 was measured in the plasma acquired from the QFT tubes including Nil, TB1, and TB2 using ELISA Development Kits (Invitrogen Inc, MN, USA), according to the manufacturer's instructions, and the results were expressed as pg/mL.

STATISTICAL ANALYSIS



- ✘ The correlation degree between IL-2 and IFN- γ was evaluated using the Spearman rank test. Test concordance between the biomarkers was calculated by kappa (k) statistic. Normal distribution of qualitative values was evaluated by the Kolmogorov–Smirnov test, and comparison of IFN γ and IL-2 secretion between the groups was performed using the Mann-Whitney U test. In addition, multivariate analysis was carried out to detect factors associated with the positive results of each test. Diagnostic accuracies of IL-2 assay and IL-2/ IFN- γ ratio were assessed by plotting the operating characteristic (ROC) curve. All statistical analyses were conducted using SPSS version 22 (IBM Corp, New York, USA). In each test, a P-value of 0.05 was considered significant

RESULTS



- ✘ A total of 237 household contacts were recruited in the final analysis, including 132 patients with LTBI, three with active TB, and 102 TB uninfected controls. The demographic and clinical features of participants are summarized in Table 1. The analysis revealed no significant differences among the groups regarding age and gender. In order to improve the accuracy of the statistical analysis, 14 patients with confirmed active TB were enrolled in addition to the three household contacts with active TB. QFT-Plus results indicated that the release of IFN- γ was significantly higher in active TB (mean IQR: 1.94 and 1.60 for TB1, and 1.94 and 1.48 for TB2) and LTBI (mean IQR: 2.46 and 3.55 for TB1, and 2.71 and 3.65 for TB2) cohorts than in the TB uninfected controls (mean IQR: 0.08 and 1.24 for TB1, and 0.07 and 1.18 for TB2). The level of produced IL-2 was significantly higher in patients with active TB (mean IQR: 180.17 and 140.97 for TB1, and 174.27 and 123.02 for TB2) and LTBI (mean IQR: 206.48 and 259.55 for TB1, and 223.84 and 267.74 for TB2) than in TB uninfected controls (mean IQR: 22.95 and 94.40 for TB1, and 23.93 and 108.77 for TB2).

- ✘ To determine the potential diagnostic value of IL-2, the cutoff point for IL-2 was calculated using the ROC curve between cases with active TB and those with LTBI. For TB1, AUC was 0.57 (95%CI: 0.46 - 0.68), and the cutoff point of 25.5 pg/mL identified active disease with 88.24% sensitivity and 36.36% specificity. Likewise, for TB2, AUC was 0.55 (95%CI: 0.55 - 0.66), and an IL-2 level of > 34.95 pg/mL predicted active disease with 88.24% sensitivity and 36.36% specificity. The obtained results also showed that IL-2 was significantly higher in LTBI cases than uninfected controls ($P = 0.001$). For TB1, the AUC was 0.83 (95%CI: 0.77 - 0.88) by ROC analysis, and the cutoff point of 14.15 showed a sensitivity of 71.21% and specificity of 90.20% for LTBI. For TB2, the AUC was 0.86 (95%CI: 0.81 - 0.91) by ROC analysis, and the cutoff point of 11.60 indicated a sensitivity of 79.55% and specificity of 89.22% for LTBI. ROC analysis for IL-2 and IL-2/IFN γ ratio in discriminating patients with the active disease from LTBI, as well as LTBI from uninfected controls, is summarized in tables 2 and 3. Finally, the study results indicated no significant differences between patients with LTBI and active TB in IL-2 serum level ($P = 0.58$).

DISCUSSION

- ✘ Cytokines play a pivotal role in the pathogenesis of TB. It is widely accepted that IFN- γ and IL-2 produced by immune cells play a pivotal role in defense against *M. tuberculosis* infection . In the last few decades, IGRAs are widely used for the indirect or immunologic diagnosis of TB infections, particularly among those at a higher risk for LTBI. However, IGRAs do not discriminate between LTBI and active disease, as the tuberculin skin test (TST)

DISCUSSION

- ✘ Although it is considered that IGRAs cannot distinguish active TB from LTBI, some authors found that the IL-2 assay, in combination with QFT-Plus, is a good candidate when is analyzed in combination. Several recent investigations indicated that IL-2-dominant functional T-cell signatures were associated with antigen clearance .

-
- ✘ The current study findings revealed that the production of IL-2 was significantly higher among cases with active TB and LTBI compared to uninfected subjects. In accordance with the current study results, several authors reported that the performance of the IL-2 assay would be considered as an alternative or adjunct diagnostic approach to M. tuberculosis infections; however, it was emphasized that IL-2 is not valuable as a stand-alone diagnostic indicator of active TB or LTBI. The ROC analysis showed that the IL-2 assay had an acceptable specificity (88.24%) in distinguishing active TB from LTBI and low sensitivity (36.36%) in diagnosis. The low sensitivity could be due to the small sample size of the active TB group compared with LTBI cases and uninfected controls.

Recent studies demonstrated the potential role of IL2 in distinguishing active TB from LTBI, suggesting a new potential biomarker . This could be due to the fact that IL-2 is differentially expressed in patients with active TB and LTBI . Bella et al., reported that CFP-10 and ESAT-6 were invaluable in distinguishing LTBI from active TB when IL-2 was measured, as for IGRAs, such as QFT and T-Spot TB test . They described that the IL-2 assay induced by AlaDH indicated relatively high sensitivity and specificity for the diagnosis of active TB.

- ✘ According to the current study analysis, in combination with the QFT-Plus test, the diagnostic sensitivity of IL-2 in distinguishing active TB from LTBI was lower than that of IL-2 alone, while the specificity of the test relatively improved. It is noteworthy that the performance of both IGRAs and IL-2 could increase the sensitivity of the test (26). It was previously described that the IL-2/IFN- γ ratio after 72 hours of stimulation with *M. tuberculosis*-specific antigens was significantly higher among patients with LTBI compared with those with active disease, with a predominant IFN- γ response

. However, the current study results did not support the previous findings. The current study findings indicated that the IL-2 assay had a relatively high specificity (90.2% for TB1 and 89.2% for TB2) in distinguishing LTBI from uninfected controls; however, it showed a relatively low sensitivity (71.2% for TB1 and 79.5% for TB2) to diagnose LTBI. In combination with the QFT-Plus test, the diagnostic sensitivity of the IL-2 assay in detecting LTBI was relatively lower than that of IL-2 alone

Table 1. Demographic and Clinical Characteristics of the Study Participants

Characteristic	Active TB	LTBI	CON	P-Value
Total No.	3	132	102	
Male, No. (%)	2 (66.6)	95 (72)	74 (72.5)	0.1
Median age (range), y	47.3 (20 - 78)	44 (11 - 63)	41 (9 - 58)	1
History of TB, No. (%)				
Yes	0 (0)	1 (0.7)	0 (0)	1
No	3 (100)	127 (96.3)	102 (100)	1
Unknown	0 (0)	4 (3)	0 (0)	
Previous TB prophylaxis or treatment, No. (%)				
Yes	0 (0)	0 (0)	1 (1)	1
No	3 (100)	128 (97)	101 (99)	1
Unknown	0 (0)	4 (3)	0 (0)	0.1

Abbreviations: LTBI, latent tuberculosis infection; CON, uninfected controls

Table 2. ROC Analysis for Cytokines in Discriminating Patients with Active TB from LTBI Cases

Cytokine	Cutoff Point (pg/mL) ^a	Sensitivity (%)	Specificity (%)	AUC (95% CI)
IL-2 in TB1 ^b	25.05	88.24	36.36	0.57 (0.46 - 0.68)
IL-2 in TB2 ^c	34.95	88.24	36.36	0.55 (0.44 - 0.66)
IL-2/IFN- γ in TB1	69.74	88.24	50.00	0.61 (0.51 - 0.72)
IL-2/IFN- γ in TB2	74.00	88.24	44.70	0.57 (0.47 - 0.67)

^aData are expressed as median concentration in pg/mL (interquartile range).

^bTB1, QFT-Plus tube 1

^cTB2, QFT-Plus tube 2

Table 3. ROC Analysis for Cytokines in Discriminating Patients with Active LTBI from Uninfected Ones

Cytokine	Cutoff Point (pg/mL) ^a	Sensitivity (%)	Specificity (%)	AUC (95% CI)
IL-2 in TB1 ^b	14.15	71.21	90.20	0.83 (0.77 - 0.88)
IL-2 in TB2 ^c	11.60	79.55	89.22	0.86 (0.81 - 0.91)
IL-2/IFN- γ in TB1	36.33	73.48	62.75	0.67 (0.60 - 0.75)
IL-2/IFN- γ in TB2	31.52	78.79	62.75	0.73 (0.66 - 0.79)

^a Data are expressed as median concentration in pg/mL (interquartile range).

^b TB1, QFT-Plus tube 1

^c TB2, QFT-Plus tube 2

- ✘ however, the specificity of IL-2/IFN- γ ratio was lower than that of the IL-2 assay. It is described that M. tuberculosis-specific antigens stimulate the IL-2 production at lower levels in LTBI cases after 24 hours; however, the amount of secreted IL-2 would be significantly higher after 72 hours in such patients compared patients with active TB and healthy controls (27). Therefore, the incubation time for IL-2 is important in obtaining valid results..

According to a study conducted by Desalegn et al., IFN- γ release increased against *M. tuberculosis*-specific antigens, but no IL-2 response was observed in HIV-LTBI coinfecting individuals with long-term HAART (28). Hence, IL2 could not be an appropriate diagnostic approach to detect TB infection among patients with HIV. Some limitations could be noted for the current study e.g, the number of participants was lower in the active TB group than LTBI and uninfected ones. The immune response was not investigated in subjects with negative QFTPlus, which might be attributed to false-negative results.

- ✘ In conclusion, the results of the current study, conducted on household contacts of cases with active TB in two high TB burden provinces of Iran, indicated that IL-2 dynamic profiles, obtained by the utilization of commercially available kits, could be used to diagnose cases with TB infection. However, the study analysis revealed that this test cannot discriminate between active TB and LTBI with acceptable sensitivity. These findings, thus, call for more prospective studies evaluating larger cohorts to determine the diagnostic value of IL-2 assay in the active TB and LTBI diagnosis.